

New insights from molecular characterization of the tick *Rhipicephalus (Boophilus) microplus* in Brazil

Novos pontos de vista sobre a caracterização molecular em carrapatos *Rhipicephalus (Boophilus) microplus*, no Brasil

Bárbara Guimarães Csordas¹; Marcos Valério Garcia^{2§}; Rodrigo Casquero Cunha³; Poliana Fernanda Giachetto⁴;
Isabella Maiumi Zaidan Blecha⁵; Renato Andreotti^{2*}

¹ Pós-graduação em Doenças Infecciosas e Parasitárias, Faculdade de Medicina, Universidade Federal de Mato Grosso do Sul – UFMS, Campo Grande, MS, Brasil

² Laboratório de Biologia Molecular do Carrapato, Departamento de Sanidade Animal, Embrapa Gado de Corte, Campo Grande, MS, Brasil

³ Programa de Pós-graduação em Biotecnologia, Universidade Federal de Pelotas – UFPel, Pelotas, RS, Brasil

⁴ Laboratório Multiusuário de Bioinformática, Embrapa Informática Agropecuária, Campinas, SP, Brasil

⁵ Programa de Pós-graduação em Ciência Animal, Faculdade de Medicina Veterinária, Universidade Federal de Mato Grosso do Sul – UFMS, Campo Grande, MS, Brasil

[§] Bolsista DCR Fundect/MS.

Received April 11, 2016

Accepted July 20, 2016

Abstract

The *Rhipicephalus (Boophilus) microplus* complex currently consists of five taxa, namely *R. australis*, *R. annulatus*, *R. (B.) microplus* clade A *sensu*, *R. microplus* clade B *sensu*, and *R. (B.) microplus* clade C *sensu*. Mitochondrial DNA-based methods help taxonomists when they are facing the morpho-taxonomic problem of distinguishing members of the *R. (B.) microplus* complex. The purpose of this study was to perform molecular characterization of ticks in all five regions of Brazil and infer their phylogenetic relationships. Molecular analysis characterized 10 haplotypes of the COX-1 gene. Molecular network analysis revealed that haplotype H-2 was the most dispersed of the studied populations (n = 11). Haplotype H-3 (n = 2) had the greatest genetic differentiation when compared to other Brazilian populations. A Bayesian phylogenetic tree of the COX-1 gene obtained strong support. In addition, it was observed that the population of *R. (B.) microplus* haplotype H-3 exhibited diverging branches among the other Brazilian populations in the study. The study concludes that the different regions of Brazil have *R. (B.) microplus* tick populations with distinct haplotypes.

Keywords: *Rhipicephalus (Boophilus) microplus*, cattle tick, COX-I gene, ITS-2 gene, Brazil.

Resumo

Carrapatos do complexo *R. (B.) microplus* se distribuem em cinco taxa: *R. australis*, *R. annulatus*, *R. (B.) microplus* clado A *sensu*, *R. microplus* clado B *sensu* e *R. (B.) microplus* clado C *sensu*. Métodos baseados no DNA mitocondrial podem auxiliar taxonomistas quando há dificuldades em estabelecer diferenças morfológicas para distinguir membros do complexo *R. (B.) microplus*. O objetivo deste estudo foi a caracterização molecular e a inferência de relações filogenéticas em carrapatos de todas as cinco regiões geográficas do Brasil. Para o gene COX-1, a análise molecular caracterizou 10 haplótipos. Na análise molecular em rede foi observado que o haplótipo H-2 é o mais disperso entre as populações (n=11). O haplótipo H-3 (n=2) foi o que obteve maior diferenciação genética ao ser comparado com outras populações brasileiras. A árvore filogenética Bayesiana de gene COX-1 gerou suporte robusto e foi observado que a população de *R. (B.) microplus* haplótipo H-3 apresentou ramificação com divergência entre as outras populações brasileiras apresentadas neste estudo. Conclui-se que as populações brasileiras possuem diversidade haplotípica com divergência entre as diversas populações de *R. (B.) microplus* no Brasil.

Palavras-chave: *Rhipicephalus (Boophilus) microplus*, carrapato-do-boi, gene COX-I, gene ITS-2, Brasil.

*Corresponding author: Renato Andreotti. Departamento de Sanidade Animal, Embrapa Gado de Corte, Avenida Rádio Maia, 830, Vila Popular, CEP 79106-550, Campo Grande, MS, Brasil. e-mail: renato.andreotti@embrapa.br

Introduction

The bovine tick *Rhipicephalus (B.) microplus* can be found in multiple tropical and subtropical regions worldwide (ESTRADA-PEÑA et al., 2006). This parasite may transmit pathogens that cause babesiosis (*Babesia bovis* and *B. bigemina*) and anaplasmosis (*Anaplasma marginale*) (PETER et al., 2005; de la FUENTE et al., 2008). Infestation of herds with *R. (B.) microplus* cause losses estimated at US\$3.24 billion per year in Brazil (GRISI et al., 2014).

In most cattle-producing countries, control methods against this tick are costly and require the use of acaricides. However, acaricide-resistant *R. (B.) microplus* populations have become a worldwide problem, and molecular ecology studies and new tick control technologies are now required to preserve cattle production (RODRÍGUEZ-VIVAS et al., 2007; ANDREOTTI et al., 2011; GUERRERO et al., 2014).

Six years ago, Labruna et al. (2009) used 12S and 16S molecular markers to show a lack of divergence among populations in the Americas, including Brazil. Later, Burger et al. (2014) used the COX-1 molecular marker to show that the *R. (B.) microplus* populations of Brazil (GenBank: KC503261) are organized in clade A, along with populations of *R. (B.) microplus* from China (GenBank: HM193863) and Cambodia (GenBank: KC503260). Thus, molecular markers may resolve very close taxonomic relationships and provide new knowledge about population structure, allowing a deeper understanding of *R. (B.) microplus* populations (BURGER et al., 2014).

Although the history of the dissemination of *R. (B.) microplus* is not well documented, the species is known to have originated in India (HOOGSTRAL, 1986). According to Barré & Uilenberg (2010), *R. (B.) microplus* originated in the southern and southeastern regions of Asia and was spread throughout the tropical and subtropical belts via cattle, arriving in Brazil between the 16th and 17th centuries.

Mitochondrial and nuclear genome markers are increasingly being used to better understand phylogenetic relationships in order to elucidate whether evolution has occurred among populations of a species (KANDUMA et al., 2012).

The DNA barcode has been proposed as a universal tool for identifying biological diversity, and can be used as a molecular marker (HEBERT et al., 2003). DNA barcoding is based on information gathered from a fragment of approximately 688 base pairs of mitochondrial DNA base sequences (mtDNA) from the cytochrome oxidase I gene (COX-I) of different species. A number of studies have shown that the DNA barcode is a universal code highly effective for the identification of species (HEBERT et al., 2004; BARRETT & HEBERT, 2005).

In addition to the DNA barcode, ribosomal DNA (rDNA) may be used as a nuclear molecular marker. The internal transcribed spacer 2 (ITS-2) is located between the 5.8S and 28S ribosomal subunits of rDNA (CRUICKSHANK, 2002). These genes are arranged in repeated units known as ribosomal cistrons, which have repeated copies and have been used to study phylogenetic

relationships (CAMPBELL et al., 1993; SONG et al., 2011; BURGER et al., 2014).

The objective of this study was to infer the phylogenetic and phylogeographic relationships among *R. (B.) microplus* tick populations in Brazil based on COX-I mitochondrial DNA gene sequences and ITS-2 nuclear DNA, and compare them with *R. (B.) microplus* tick populations from other countries.

Materials and Methods

Tick collection

Ticks were obtained from 22 locations in different geographical regions of Brazil (Figure 1, Table 1). A pool of larvae and 22 engorged females were collected from natural environments. The morphology and identification key was based on Barros-Battesti et al. (2006). Samples were stored in an ultra-freezer at -80°C for subsequent analysis.

DNA extraction and PCR amplification

DNA extraction was performed on a pool of larvae and individual samples of engorged females using DNAzol® Reagent (Invitrogen) according to the manufacturer's recommendations. The DNA was quantified using a Nano Drop 1000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA), and an A260nm/A280nm ratio above 1.6 was established as acceptable. The sample concentration was corrected to $50\text{ ng }\mu\text{L}^{-1}$. Next, the DNA was used in a polymerase chain reaction (PCR) for COX-I and ITS-2 region amplification with new primers designed by our group. PCR generated products of 643bp and 580bp for COX-1 and ITS-2, respectively.

COX-I gene fragments were amplified using the COX-I.2F (5'-CTTCAGCCATTTTACCGCGA-3') and COX-I.2R (5'-CTCCGCCTGAAGGGTCAAA-3') starter oligonucleotides. ITS-2 fragments were amplified with the ITS-2F (5'-CGGATCACATATCAAGAGAG-3') and ITS-2R (5'-CCCAACTGGAGTGGCCCAGTTT-3') primers. The PCR was standardized for a final volume of 25 μL with 1X buffer, 1.5 mM MgCl₂, 25 mM dNTPs, 10 μM of each primer, 0.5 unit of Platinum® Taq DNA Polymerase High Fidelity (Invitrogen by Life Technologies™, Massachusetts, USA), and 1 μL of DNA at $50\text{ ng }\mu\text{L}^{-1}$. PCR conditions were optimized for each reaction, and the annealing temperature was adjusted to suit the primers used. General PCR conditions were 94°C for 2 min, followed by 35 cycles of 94°C for 1 min, $56^{\circ}\text{C}/60^{\circ}\text{C}$ (COX-I/ITS-2, respectively) for 30s, 68°C for 1 min, and a final extension of 68°C for 3 min. PCR products were visualized in a 2% agarose gel stained with ethidium bromide. The PCR-amplified products were purified using a Purelink® kit (Invitrogen by Life Technologies™, Massachusetts, USA). Fragments were then cloned with pGEM®-T Easy Vector Systems cloning vector according to the manufacturer's manual (Promega, Madison, WI-USA) and sequenced as described below.




Figure 1. Geographic locations where *Rhipicephalus (Boophilus) microplus* ticks were collected in five regions of Brazil.

Genetic analysis

All analyses generated in this study were performed *in silico*. Sanger sequencing was performed in a 48 well plate with robotic instrumentation using T7 Transcription Start (5'-CTAATACGACTCACTATAGGG-3') universal sequencing primer (Promega, Madison, WI-USA). Samples were sequenced using an Applied Biosystems™ ABI 3730 DNA Analyzer and the conditions described by Song et al. (2011). The sequencing reactions were done using the Big Dye® Terminator v3.1 cycle sequencing kit. The runs were performed in 36-cm capillaries using the POP7 polymer, and the sequences were generated by the Sequencing Analysis Software v5.3.1 through the Caller KB Phred program (EWING & GREEN, 1998; EWING et al., 1998).

Plasmid sequences were identified and removed using an NCBI (VecScreen) tool (ALTSCHUL et al., 1997), and a hash search algorithm was used to remove the contaminant sequences.

Consensus sequences were aligned with the GenBank data using the BLASTN program (ALTSCHUL et al., 1997).

The partition homogeneity test was conducted using the ARLEQUIN program, version 3.5.1.2 (EXCOFFIER & LISCHER, 2010). The aligned COX-I and ITS-2 sequences comprised 681bps and 649bps, respectively. Database sequence *R. (B.) microplus* MS- Brazil (GenBank: KC503261) clade *A sensu* Burger et al. (2014), was used for alignment with samples from this study. This sequence alignment revealed 10 haplotypes (H1-H10) based on COX-I genes alone (Figure 2). Distinct sequences of COX-I (KP226159-KP226180) and ITS-2 (KP226139-KP226158) were deposited in NCBI GenBank.

A median-joining analysis implemented in the program Network Version 5.0 (BANDELTA et al., 1999) was used for the intraspecific analysis of the evolutionary relationships among haplotypes. Uncorrected (p) pairwise genetic distances were calculated using ARLEQUIN program, version 3.5.1.2

Table 1. Brazilian states, geographic coordinates related to the collection and GenBank accession number of this study.

Region	State	Geographic coordinates	GenBank	GenBank
			(COX-1)	(ITS2)
North	AC	9°59'30"S/67°48'36"W	KP226159	KP226139
	AM	3°6'0"S/60°1'0"W	KP226162	-
	PA	1°17'49"S/47°55'19"W	KP226163	KP226153
	RO	8°45'43"S/63°54'14"W	KP226178	KP226142
	RR	2°49'10"S/60°40'17"W	KP226161	-
	TO	10°11'4"S/48°20'01"W	KP226180	KP226146
	MA	4°51'32"S/43°21'21"W	KP226160	KP226157
Northeast	CE	3°43'6"S/38°32'34"W	KP226177	KP226140
	RN	5°47'42"S/35°12'32"W	KP226172	KP226145
	PB	6°57'46"S/35°41'31"W	KP226176	KP226141
	PI	9°4'26"S/44°21'32"W	KP226174	KP226148
	BA	12°58'16"S/38°30'39"W	KP226168	KP226156
Midwest	MT	13°49'44"S/56°04'56"W	KP226167	KP226143
	GO	15°47'56"S/47°52'00"W	KP226166	KP226151
	MS	20°26'34"S/54°38'45"W	KP226164	KP226155
	ES	20°45'50"S/41°31'58"W	KP226175	KP226154
Southeast	MG	18°55'8"S/48°16'37"W	KP226171	KP226150
	RJ	21°38'24"S/41°3'3"W	KP226165	KP226149
	SP	21°15'18"S/48°19'19"W	KP226170	KP226158
	PR	25°23'42"S/51°27'28"W	KP226179	KP226147
South	SC	27°20'27"S/51°37'4"W	KP226173	KP226144
	RS	31°46'19"S/52°20'34"W	KP226169	KP226152

(EXCOFFIER & LISCHER, 2010) to assess the genetic divergence of *R. (B.) microplus* in both COX-I and ITS-2 genes. To assess the level of genetic differentiation, we used the program DnaSP 5.0 (LIBRADO & ROZAS, 2009). Haplotype diversity (Hd), nucleotide diversity (pi), number of haplotypes (h), and genetic differentiation (FST) were determined.

The levels of genetic differentiation were defined as $F_{ST} > 0.25$ (great differentiation), 0.15 to 0.25 (moderate differentiation) and $F_{ST} < 0.05$ (negligible differentiation) (WRIGHT, 1931). The levels of gene flow were defined as $Nm > 1$ (high gene flow), 0.25 to 0.99 (intermediate gene flow) and $Nm < 0.25$ (low gene flow). The distinct COX-I and 16S rRNA haplotypes identified in the Brazilian *R. (B.) microplus* ticks were aligned with all representative sequences of Rhipicephalus taxa available in the NCBI GenBank. *R. annulatus* was used as an outgroup for the construction of Bayesian phylogenetic trees based on Brazilian *R. (B.) microplus* ticks COX-I only; *Dermacentor nitens* was used as an outgroup for the construction of Bayesian phylogenetic trees based on COX-I and ITS-2 sequences. The Bayesian phylogenetic analysis performed using MrBayes 3.1 (HUELSENBECK & RONQUIST, 2001; RONQUIST & HUELSENBECK, 2003) were plotted using FigTree software version 1.4.2 (TREE BIO, 2016).

The ARLEQUIN software revealed population structure by means of AMOVA, which uncovers the existence of population differentiation at both intra- and inter-population levels (EXCOFFIER & LISCHER, 2010).

Results and Discussion

Nucleotide analyses and Haplotype

The phylogenetic and phylogeographic relations of the Brazilian *R. (B.) microplus* were inferred from the COX-I sequences (22 samples), *R. (B.) microplus* Brazil (GenBank: KC503261) and the ITS2 sequences (20 samples) obtained from all five Brazilian geographic regions (Figure 1).

This is the only study that has compared a representative sample of *R. (B.) microplus*, which together represent the whole country and the authors were careful to represent each region by just having differentiated biogeography. Assays performed by Dantas-Torres (2015), can demonstrate that certain factors such as climate change and biodiversity boost the expansion of tick populations.

The COX-I dataset showed haplotype diversity (0.767) greater than that revealed by the ITS-2 (0.567) gene. However, the ITS-2 showed greater nucleotide diversity (0.00277) than the COX-I (0.00432) dataset. Haplotype network analysis (COX-I) revealed ten distinct haplotype clusters among Brazilian populations, with no clear separation by tick populations/geographical areas, indicating that an overlap of the ten genetically divergent haplotypes of *R. (B.) microplus* (Figure 2A) exists.

Haplotype H-2 was the most widespread haplotype (n = 11) in Northern, Midwest, Southeast and Southern populations (Figure 2B). Haplotypes H-5, H-6 and H-8 (Northern region) originates from H-4 and H-2 haplotypes, as shown in Figure 2B.




Figure 2. (A) Bayesian phylogenetic tree recovered for cytochrome oxidase subunit 1 gene (COX-I) of 643 nucleotides. *R. annulatus* (GenBank: KC503256) was used as the outgroup (B) median joining network of *R. (B.) microplus* from five different regions populations in Brazil. A circle represents each haplotype. Relative areas of the circles indicate haplotype frequency of each sample.

The ITS-2 Bayesian phylogenetic tree revealed one *R. (B.) microplus* genetic clade (Figure 4). The COX-I Bayesian phylogenetic tree revealed two genetic clades of *R. (B.) microplus*, Clade A (Brazil, Texas, Malaysia, *R. australis*) and Clade B (*R. (B.) microplus* China and *R. annulatus*). *Rhipicephalus (B.) microplus* clade A showed a sister relationship with *R. australis*. By contrast, *R. (B.) microplus* clade B is more closely related to *R. annulatus*, with which it forms a sister group relationship, corroborating the data by Burger et al. (2014) (Figure 3).

Table 4. Analysis of molecular variance of molecular marker, gene COX-1.

Source of variation	d.f	Sum of squares	Variance Components	Percentage of variation
Among populations	9	25.059	1.34 _{Va}	82.69
Within populations	13	3.636	0.28 _{Vb}	17.31
Total	22	28.696	1.62	
Fixation Index FST:	0.8269	P < 0.001		

The *R. (B.) microplus* populations in Brazil have internal nodes, indicating the occurrence of subpopulations (GenBank: KP226160, KP226167, KP226169, KP226172, KP226174, KP226177). The haplotype H-3 formed another smaller node (GenBank: KP226176, KP226178) (Figure 3).

The intraspecific genetic distance or haplotype frequency, like that of haplotype H2 reported here, are notably higher than previously described for *R. (B.) microplus sensu* Low et al. (2015). Furthermore, the two, and perhaps three, genetic assemblages inferred from Brazilian *R. (B.) microplus* are more genetically diverse than those reported from other regions (Figure 3).

This study sought to obtain high support for the topology of tree and the COX-I molecular marker was found to have the advantage of presenting greater mutation rates than the nuclear marker, corroborating the data presented by Lv et al. (2014). However, the ITS-2 molecular marker failed to resolve discrepancies between the populations of *R. (B.) microplus* and this study corroborates the findings of Burger et al. (2014) in which they showed that the ITS-2 molecular marker has insufficient power to distinguish between very close species.

The comparative analysis in this study shows that *R. (B.) microplus* ticks represent at least two different populations. Moreover, this study also attempted to investigate whether haplotypes are




Figure 3. Bayesian phylogenetic tree recovered for cytochrome oxidase subunit 1 gene (COX-I) of 643 nucleotides, for *Rhipicephalus* taxa samples. In blue and green are the smallest nodes formed the Brazilian populations of *R. (B.) microplus*. *Dermacentor nitens* (GenBank: KC503258) was used as the outgroup.

be able to show how these populations differ in their inter- and intra- relationships.

Acknowledgements

We thank FUNDECT - MS, the State of Mato Grosso do Sul, CAPES and CNPq for financial support.

References

- Altschul SE, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997; 25(17): 3389-3402. <http://dx.doi.org/10.1093/nar/25.17.3389>. PMID:9254694.
- Andreotti R, Guerrero FD, Soares MA, Barros JC, Miller RJ, Léon AP. Acaricide resistance of *Rhipicephalus (Boophilus) microplus* in State of Mato Grosso do Sul, Brazil. *Rev Bras Parasitol Vet* 2011; 20(2): 127-133. <http://dx.doi.org/10.1590/S1984-29612011000200007>. PMID:21722487.
- Bandelt HJ, Forster P, Röhl A. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* 1999; 16(1): 37-48. <http://dx.doi.org/10.1093/oxfordjournals.molbev.a026036>. PMID:10331250.
- Barré N, Uilenberg G. Spread of parasites transported with their hosts: case study of two species of cattle tick. *Rev Sci Tech* 2010; 29(1): 149-160, 135-147. PMID:20617654.
- Barros-Battesti DM, Arzua M, Bechara GH. *Carrapatos de importância médico-veterinária da região neotropical: Um guia ilustrado para identificação de espécies*. São Paulo: Instituto Butantan, 2006. 223 p.
- Barrett RDH, Hebert PDN. Identifying spiders through DNA barcodes. *Can J Zool* 2005; 83(3): 481-491. <http://dx.doi.org/10.1139/z05-024>.
- Busch JD, Stone NE, Nottingham R, Araya-Anchetta A, Lewis J, Hochhalter C, et al. Widespread movement of invasive cattle fever ticks (*Rhipicephalus microplus*) in southern Texas leads to shared local infestations on cattle and deer. *Parasit Vectors* 2014; 7(1): 188. <http://dx.doi.org/10.1186/1756-3305-7-188>. PMID:24742041.
- Burger TD, Shao R, Barker SC. Phylogenetic analysis of mitochondrial genome sequences indicates that the cattle tick, *Rhipicephalus (Boophilus) microplus*, contains a cryptic species. *Mol Phylogenet Evol* 2014; 76: 241-253. <http://dx.doi.org/10.1016/j.ympev.2014.03.017>. PMID:24685498.
- Campbell BC, Steffen-Campbell JD, Werren JH. Phylogeny of the *Nasonia* species complex (Hymenoptera: Pteromalidae) inferred from an internal transcribed spacer (ITS2) and 28S rDNA sequences. *Insect Mol Biol* 1993; 2(4): 225-237. <http://dx.doi.org/10.1111/j.1365-2583.1994.tb00142.x>. PMID:9087560.
- Cruikshank RH. Molecular markers for the phylogenetics of mites and ticks. *Syst Appl Acarol* 2002; 7(1): 3-14. <http://dx.doi.org/10.11158/saa.7.1.1>.
- Dantas-Torres F. Climate change, biodiversity, ticks and tick-borne diseases: the butterfly effect. *Int J Parasitol Parasites Wildl* 2015; 4(3): 452-461. <http://dx.doi.org/10.1016/j.ijppaw.2015.07.001>. PMID:26835253.
- de la Fuente J, Estrada-Peña A, Venzal JM, Kocan KM, Sonenshine DE. Overview: ticks as vectors of pathogens that cause disease in humans and animals. *Front Biosci* 2008; 13(13): 6938-6946. <http://dx.doi.org/10.2741/3200>. PMID:18508706.
- Estrada-Peña A, Bouattour A, Camicas JL, Guglielmone A, Horak I, Jongejan F, et al. The known distribution and ecological preferences of the tick subgenus *Boophilus* (Acari: Ixodidae) in Africa and Latin America. *Exp Appl Acarol* 2006; 38(2-3): 219-235. <http://dx.doi.org/10.1007/s10493-006-0003-5>. PMID:16596355.
- Ewing B, Green P. Base-calling of automated sequencer traces using *phred*. II. Error probabilities. *Genome Res* 1998; 8(3): 186-194. <http://dx.doi.org/10.1101/gr.8.3.186>. PMID:9521922.
- Ewing B, Hillier L, Wendl MC, Green P. Base-calling of automated sequencer traces using *phred*. I. Accuracy assessment. *Genome Res* 1998; 8(3): 175-185. <http://dx.doi.org/10.1101/gr.8.3.175>. PMID:9521921.
- Excoffier L, Lischer HEL. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 2010; 10(3): 564-567. <http://dx.doi.org/10.1111/j.1755-0998.2010.02847.x>. PMID:21565059.
- Grisi L, Leite RC, Martins JR, Barros AT, Andreotti R, Caçado PH, et al. Reassessment of the potential economic impact of cattle parasites in Brazil. *Rev Bras Parasitol Vet* 2014; 23(2): 150-156. <http://dx.doi.org/10.1590/S1984-29612014042>. PMID:25054492.
- Guerrero FD, Andreotti R, Bendele KG, Cunha RC, Miller RJ, Yeater K, et al. *Rhipicephalus (Boophilus) microplus* aquaporin as an effective vaccine antigen to protect against cattle tick infestations. *Parasit Vectors* 2014; 7: 475. PMID:25306139.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR. Biological identifications through DNA barcodes. *Proc Biol Sci* 2003; 270(1512): 313-321.
- Hebert PDN, Stoeckle MY, Zemplak TS, Francis CM. Identification of birds through DNA barcodes. *PLoS Biol* 2004; 2(10): e312. <http://dx.doi.org/10.1371/journal.pbio.0020312>. PMID:15455034.
- Hoogstral H. Theobald Smith: his scientific work and impact. *Bull Entomol Soc Am* 1986; 32(1): 22-35. <http://dx.doi.org/10.1093/besa/32.1.22>.
- Huelsenbeck JP, Ronquist F. MRBAYES: bayesian inference of phylogenetic trees. *Bioinformatics* 2001; 17(8): 754-755. <http://dx.doi.org/10.1093/bioinformatics/17.8.754>. PMID:11524383.
- Ibelli AM, Ribeiro AR, Giglioti R, Regitano LC, Alencar MM, Chagas ACS, et al. Resistance of cattle of various genetic groups to the tick *Rhipicephalus microplus* and the relationship with coat traits. *Vet Parasitol* 2012; 186(3-4): 425-430. <http://dx.doi.org/10.1016/j.vetpar.2011.11.019>. PMID:22115946.
- Kanduma EG, Mwacharo JM, Sunter JD, Nzuki I, Mwaura S, Kinyanjui PW, et al. Micro- and minisatellite-expressed sequence tag (EST) markers discriminate between populations of *Rhipicephalus appendiculatus*. *Ticks Tick Borne Dis* 2012; 3(3): 128-136. <http://dx.doi.org/10.1016/j.ttbdis.2012.05.001>. PMID:22789728.
- Labruna MB, Naranjo V, Mangold AJ, Thompson C, Estrada-Peña A, Guglielmone AA, et al. Allopatric speciation in ticks: genetic and reproductive divergence between geographic strains of *Rhipicephalus (Boophilus) microplus*. *BMC Evol Biol* 2009; 9(1): 46. <http://dx.doi.org/10.1186/1471-2148-9-46>. PMID:19243585.
- Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 2009; 25(11): 1451-1452. <http://dx.doi.org/10.1093/bioinformatics/btp187>. PMID:19346325.
- Low VL, Tay ST, Kho KL, Koh FX, Tan TK, Lim YAL, et al. Molecular characterisation of the tick *Rhipicephalus microplus* in Malaysia: new insights into the cryptic diversity and distinct genetic assemblages throughout the world. *Parasit Vectors* 2015; 8(1): 341. <http://dx.doi.org/10.1186/s13071-015-0956-5>. PMID:26104478.

Lv J, Wu S, Zhang Y, Chen Y, Feng C, Yuan X, et al. Assessment of four DNA fragments (COI, 16S rDNA, ITS2, 12S rDNA) for species identification of the Ixodida (Acari: Ixodida). *Parasit Vectors* 2014; 7(1): 93. <http://dx.doi.org/10.1186/1756-3305-7-93>. PMID:24589289.

Peter RJ, Van den Bossche P, Penzhorn BL, Sharp B. Tick, fly, and mosquito Control-Lessons from the past, solutions for the future. *Vet Parasitol* 2005; 132(3-4): 205-215. <http://dx.doi.org/10.1016/j.vetpar.2005.07.004>. PMID:16099104.

Rodríguez-Vivas RI, Rivas AL, Chowell G, Fragoso SH, Rosario CR, García Z, et al. Spatial distribution of acaricide profiles (*Boophilus microplus* strains susceptible or resistant to acaricides) in southeastern Mexico. *Vet Parasitol* 2007; 146(1-2): 158-169. <http://dx.doi.org/10.1016/j.vetpar.2007.01.016>. PMID:17349747.

Ronquist F, Huelsenbeck JP. MRBAYES 3: bayesian phylogenetic inference under mixed models. *Bioinformatics* 2003; 19(12): 1572-1574. <http://dx.doi.org/10.1093/bioinformatics/btg180>. PMID:12912839.

Song S, Shao R, Atwell R, Barker S, Vankan D. Phylogenetic and phylogeographic relationships in *Ixodes holocyclus* and *Ixodes cornuatus* (Acari: Ixodidae) inferred from COX1 and ITS2 sequences. *Int J Parasitol* 2011; 41(8): 871-880. <http://dx.doi.org/10.1016/j.ijpara.2011.03.008>. PMID:21540032.

Tree Bio. *FigTree* [online]. London; 2016 [cited 2016 Apr 11]. Available from: <http://tree.bio.ed.ac.uk/>

Wright S. Evolution in mendelian populations. *Genetics* 1931; 16(2): 97-159. PMID:17246615.